



Protocol Booklet

Product Code(s)	HB18560, HB19791, HB16587, HB19874
Product Name(s)	Biotin Tyramide Signal Amplification Kit – AF488 Biotin Tyramide Signal Amplification Kit – Janelia Fluor® 525 Biotin Tyramide Signal Amplification Kit – Janelia Fluor® 549 Biotin Tyramide Signal Amplification Kit – Janelia Fluor® 646
Purpose	Fluorescent imaging of low abundance targets through tyramide amplification imaging.

Please note: This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use



Contents

Product Overview	3
Components & Storage	4
Protocol	4
Preparing reagents and advice on optimization	4
Safety	5
Immunohistochemistry – Free Floating Protocol	5
Immunocytochemistry Protocol	6
Guidelines, precautions, troubleshooting	7
Contact	7
For customers in the UK, Europe and Rest of the World	7
For customers in the USA, Canada and South America	7

Product Overview

Tyramide Signal Amplification (TSA) is a technique used to amplify signals in immunofluorescence. TSA is a highly sensitive method enabling detection of low abundance proteins. TSA can easily be added into the workflow of IHC and ICC. TSA works by using a HRP conjugated secondary antibody which uses biotin tyramide as a substrate in the presence of hydrogen peroxide. Tyramide becomes activated and covalently binds to neighboring tyrosine residues, creating a dense biotin labelled region in the vicinity of the primary antibody. Fluorescently conjugated streptavidin binds with high affinity to biotin, enabling detection through immunofluorescence. TSA is also compatible with multiplexing with fluorescent secondary antibodies providing that the other antibodies used are from different species.




Figure 1: Representative images of Tyramide Signal Amplification compared to detection using a biotin secondary antibody with fluorescent streptavidin. A TSA of parvalbumin in the rat hippocampus using TSA kit – AF488 (HB16462). **B** Detection of parvalbumin using a biotin secondary antibody and Streptavidin AF488. Images were taken under identical conditions.




Figure 2: Diagram of Tyramide Signal Amplification. TSA uses a secondary HRP antibody which converts Biotin Tyramide from its inactive form to activated form in the presence of hydrogen peroxide. Activated Biotin Tyramide can covalently bind to tyrosine side chains on neighboring proteins. Fluorescent streptavidin can be used for detection of biotin, resulting in amplified signal.



Components & Storage

This kit contains:

- DMSO (100µl)
- Biotin tyramide (1mg)
- Lyophilized streptavidin fluorophore conjugate (0.1mg)
- Stop solution (100ml)
- DAPI (100µl or 1mg/ml)

Note: Please store all kit components at -20°C. Please let all components warm to room temperature before use and also note that biotin tyramide and the streptavidin fluorophore conjugate require reconstitution before use. Dissolve the biotin tyramide (1mg) in DMSO (100 µl) to make a stock solution of 10 mg/ml. Aliquot and store at -20°C. Reconstitute the Streptavidin with 100 µL ddH₂O to make a stock concentration of 1 mg/ml. Store at 4°C for short term storage, or add 50% glycerol and store at -20°C for long term storage.

This kit additionally requires:

- Sodium borohydride (NaBH₄) (for IHC)
- PBS
- Tween® 20
- Triton™ X-100 (for ICC)
- Serum (for blocking)
- Bovine serum albumin (BSA)
- Glycine (for IHC)
- H₂O₂
- Primary antibody
- HRP conjugated secondary antibody
- (Optional) Avidin biotin blocking

Protocol

Preparing reagents and advice on optimization

Before starting experiments it is worth preparing all the necessary buffers in advance:

- **PBST:** 0.1% Tween20 in PBS
- **Exogenous peroxidase quenching buffer:** 0.3% H₂O₂ in PBS
- **Blocking buffer:** 2% BSA and 3% goat serum in PBS
- **Amplification buffer:** 1-10 µg/mL biotin tyramide (this may require optimization – please see below), 0.1% Tween20, 0.003% hydrogen peroxide in PBS

It is likely that parameters will need optimizing for your specific experimental conditions. This includes:

- Concentration of primary antibody (less antibody may be required due to the amplification process, depending on the strength of the signal)
- Concentration of HRP secondary antibody
- Concentration of Biotin Tyramide (1-10 µg/mL is recommended)
- Biotin Tyramide incubation time (2-30 minutes is recommended, cells are likely to need less time than tissue)
- Concentration of fluorescent streptavidin for detection

The concentrations of biotin tyramide and the timings given in this protocol were used for amplification of the signal from antibodies staining parvalbumin ([HB6457](#)) in free floating rat brain sections and GluA₁₋₄ (see [Nusser et al., 1998](#)) in neuronal cell culture. It is recommended that titrations are performed to determine the optimal conditions for TSA.

If excess background signal is observed this can be due to endogenous biotin in the sample. It is possible to use kits which involve incubation with avidin (to bind biotin in the tissue), followed by incubation of biotin (to bind all 4 biotin binding sites on avidin). This step should be carried out after blocking the sample.



To be sure that the reaction is not occurring non-specifically, it is recommended to carry out a control in which no primary antibody is present. Any fluorescence observed in this sample would suggest an artefact of the enzymatic reaction is being observed. Further optimization of the TSA conditions as well as blocking endogenous biotin can eliminate non-specific signal.

Safety

Many of the chemicals used in immunocytochemistry and immunohistochemistry have dangerous properties and can cause serious harm if not handled correctly. Always follow local rules and read the full COSHH document for any chemical that you have not used previously. Always wear appropriate PPE such as a lab coat and gloves. Specifically highlighted hazards:

- Formaldehyde is a highly toxic poison, skin sensitizer and carcinogen and should only ever be used in a fume hood with full PPE.
- Sodium borohydride is toxic, causes severe skin burns and eye damage in addition to reacting rigorously with H₂O to create flammable gas which can ignite spontaneously.
- Hydrogen peroxide can cause irritation to the eyes, nose skin and throat.

Immunohistochemistry – Free Floating Protocol

For more information on immunohistochemistry see our [immunohistochemistry protocol](#). 1 mL of kit components is used per well in a 12 well dish.

1. Perform any [antigen retrieval steps](#) as required for primary antibodies to bind.
2. Wash sections three times in PBS-Tween for 5 minutes per wash.
3. Incubate slides in 1% NaBH₄ in PBS-Tween for 15 minutes, RT.
4. Incubate sections in **exogenous peroxidase quenching buffer** (0.3% H₂O₂ in PBS) for 15 minutes at RT.
5. Wash sections three times with PBST for 5 minutes per wash.
6. Incubate sections in **blocking buffer** (2% BSA, 3% goat serum in PBST) for 1 hour at RT.
7. (Optional) To reduce background staining an endogenous biotin blocking kit can be used at this point.
8. Dilute primary antibody in **blocking buffer** to the recommended concentration. Incubate sections at 4°C overnight on a rocker.
9. Wash sections three times with PBST for 5 minutes per wash.
10. Dilute the HRP secondary antibody to 4 µg/mL in **blocking buffer**. Incubate sections in HRP conjugated secondary for 1 hour at RT.
11. Wash sections three times with PBST for 5 minutes per wash.
12. Incubate sections in **amplification buffer** (1 mL PBST containing 0.003% H₂O₂, 1 µL of 10 µg/mL Biotin Tyramide, or desired concentration) for 15 minutes at RT.
13. Incubate sections with 1 mL stop solution for 10 minutes at RT to quench the reaction.
14. Wash sections three times with PBST for 5 minutes per wash.
15. Complete steps from this point in the dark. Incubate sections with fluorescent streptavidin (1 µL of 1 mg/mL fluorescent streptavidin in 1 mL blocking buffer) for 2 hours at RT.
16. Wash sections three times with PBST for 5 minutes per wash.
17. Stain sections with DAPI (1µL of 1 mg/mL DAPI in 1 mL PBS) for 10 minutes at RT.



18. Wash sections with PBST for 5 minutes and then wash sections in ddH₂O.
19. Mount sections onto a slide using a paintbrush. Once dry, add a drop or two of [mounting media](#) and add a coverslip, taking care to avoid bubbles.
20. Once the mounting media is dry store slides at 4°C in the dark until ready to image.

Immunocytochemistry Protocol

For more information on immunocytochemistry see our [immunocytochemistry protocol](#). 1 mL of kit components is used per well in a 12 well dish.

1. Permeabilize coverslips by incubating in PBS containing 0.1% Triton™ X-100 for 10 minutes at RT.
2. Incubate coverslips in **exogenous peroxidase quenching buffer** (0.3% H₂O₂ in PBS) for 15 minutes at RT.
3. Wash coverslips three times with PBST for 5 minutes per wash.
4. Incubate coverslips in **blocking buffer** (2% BSA, 3% goat serum in PBST) for 1 hour at RT.
5. (Optional) To reduce background staining an endogenous biotin blocking kit can be used at this point.
6. Dilute primary antibody in **blocking buffer** to recommended concentration. Incubate coverslips at 4°C overnight.
7. Wash coverslips three times with PBST for 5 minutes per wash.
8. Dilute HRP conjugated secondary antibody to 1.67 ug/mL (or desired concentration) in **blocking buffer** for 1 hour at RT.
9. Wash coverslips three times with PBST for 5 minutes per wash.
21. Incubate coverslips in **amplification buffer** (containing 0.003% H₂O₂, 1 µg/mL Biotin Tyramide, or desired concentration, in PBST) for 2 minutes at RT.
10. Incubate coverslips with stop solution for 10 minutes at RT to quench the reaction.
11. Wash coverslips three times with PBST for 5 minutes per wash.
12. Incubate coverslips with fluorescent streptavidin (1 µL of 1 mg/mL fluorescent streptavidin in 1 mL **blocking buffer**) for 1 hour at RT.
13. Wash coverslips three times with PBST for 5 minutes per wash.
14. Stain sections with DAPI (1µL of 1 mg/mL DAPI in 1 mL PBS) for 2 minutes at RT.
15. Wash coverslips with PBST for 5 minutes and then wash sections in ddH₂O.
16. Mount coverslips onto a slide using a drop of [mounting media](#), taking care to avoid bubbles.
17. Once the mounting media is dry store slides at 4°C in the dark until ready to image.



Guidelines, precautions, troubleshooting

Please follow the below table to resolve any problems encountered when using this kit. For any problems not listed or for any further advice please contact our technical support team at technicalhelp@helloworldbio.com.

Problem	Potential Cause
High background	Too long incubation with biotin tyramide. Try reducing the incubation time and repeating the experiment.
	Too high concentration of HRP antibody or biotin tyramide. Try repeating the experiment with reduced concentrations or either or both.
	High levels of endogenous biotin in the sample. Try using a avidin biotin blocking kit.
	Exogenous peroxidases have not been quenched. Ensure incubation of sample with 0.3% H ₂ O ₂ for 10 minutes prior to labelling with antibodies.
Low signal	Too little primary antibody is present. Try increasing the primary antibody concentration
	Too little HRP antibody is present. Try increasing the HRP antibody incubation. Ensure the HRP antibody binds to the correct species for your primary antibody.

Observe safe laboratory practice and consult the safety datasheet. Please see the datasheet on our website for general guidelines, precautions, limitations on the use of the assay kit.

Contact

For customers in the UK, Europe and Rest of the World

Customer Care customercare@helloworldbio.com

Technical support technicalhelp@helloworldbio.com

By telephone: +44(0)117 318 0505

By fax: +44(0)117 981 1601

Opening hours: 8.30 am - 5.00 pm GMT weekdays

For customers in the USA, Canada and South America

Customer Care customercare-usa@helloworldbio.com

Technical support technicalhelp@helloworldbio.com

By telephone: +1-609-683-7500

By fax: +1-609-228-4994

Opening hours: 9.00 am - 5.00 pm EST weekdays