

Hello Bio, Inc.
304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500
F. 609-228-4994

customercare-usa@m2stage.hellobio.com



DATASHEET

SR 95531 hydrobromide (Gabazine)

Product overview

Name	SR 95531 hydrobromide (Gabazine)
Cat No	HB0901
Alternative names	Gabazine GBZ
Biological action	Antagonist
Purity	>98%
Customer comments	<p><i>We regularly use Hello Bio Gabazine (SR 95531) in the lab. We especially like the formulation where you only need to add 1ml of water to make a 10mM stock solution. Verified customer, The University of Newcastle</i></p>


I am satisfied with the quality, quick delivery and follow-up of your product. Verified customer, Shimane University

We used our first aliquot of SR95531 (Gabazine) last week. The experiment was a critical one for us and the SR95531 worked exactly as expected – 100% block of a GABAergic IPSP (inhibitory postsynaptic potential). Verified customer, University of Michigan

Good compound. This compound is routinely used in our lab to isolate AMPA and NMDA currents. So we are using it a lot every day. There are no complaints about it! Verified customer, Karolinska Institutet

Description	Selective, competitive GABA _A receptor antagonist
-------------	--

Images



Biological Data

Biological description

SR 95531 hydrobromide (Gabazine) is a selective and competitive GABA_A receptor antagonist (K_i = 150 nM for displacement of [³H]-GABA from rat membranes).

SR 95531 (Gabazine) displaces GABA from the GABA_AR agonist binding site to prevent receptor activation. It also acts as a negative allosteric inhibitor of channel opening to inhibit GABA_A receptor activation by anaesthetic agents. It also displays low affinity glycine receptor inhibition.

SR 95531 (Gabazine) inhibits GABA-induced Cl⁻ currents to reduce GABA-mediated synaptic inhibition.

SR 95531 additionally shows convulsive actions.

Application notes

Gabazine (SR 95531) is commonly used to reduce levels of inhibition by antagonising GABA_A receptors. It is commonly used at concentrations between 10 – 200 μM.

Gabazine (SR 95531) from Hello Bio blocks spontaneous inhibitory post synaptic currents (IPSC) and evoked IPSCs (see Fig 1 above). It was effective at 1 μM and completely blocked GABA_A receptors at 20 μM.

#Protocol 1: Evoked and spontaneous inhibitory post synaptic currents (IPSCs)

- Whole cell voltage clamp recordings were obtained from layer V neurons of the mouse prelimbic cortex brain slice.
- A stimulating electrode was placed in layers II/III and IPSCs were evoked by a single square (150 μs) pulse every 10 sec at a stimulus intensity that gave a reliable IPSC.
- IPSCs were evoked at a range of neuron holding voltages to measure the reversal potential of the current to ensure it was GABAergic.
- Neurons were held at 0mV and IPSCs continuously stimulated and recorded in response to 5 min applications of varying concentrations of Gabazine until complete receptor inhibition.
- Spontaneous IPSCs were recorded before and after addition of Gabazine by holding the neuron at 0mV and recording for 10 sec.
- All recordings for IPSCs were made in the presence of AMPAR antagonists.

Solubility & Handling

Storage instructions

Room temperature

Solubility overview

Soluble in water (25mM) and in DMSO (100mM)

Important

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use.

Chemical Data


Chemical name

6-Imino-3-(4-methoxyphenyl)-1(6H)-pyridazinebutanoic acid hydrobromide

Molecular Weight

368.23

Chemical structure



Molecular Formula

C₁₅H₁₇N₃O₃.HBr

CAS Number

104104-50-9

PubChem identifier

107895

SMILES

COC1=CC=C(C=C1)C2=NN(C(=N)C=C2)CCCC(=O)O.Br

Source

Synthetic

InChi

InChI=1S/C15H17N3O3.BrH/c1-21-12-6-4-11(5-7-12)13-8-9-14(16)18(17-13)10-2-3-15(19)20;/h4-9,16H,2-3,10H2,1H3,(H,19,20);1H

InChiKey

GFZHNFOGCMEYTA-UHFFFAOYSA-N

MDL number

MFCD00055135

Chemical name	6-Imino-3-(4-methoxyphenyl)-1(6 <i>H</i>)-pyridazinebutanoic acid hydrobromide
Appearance	White solid

References

Biochemical characterization of the interaction of three pyridazinyl-GABA derivatives with the GABAA receptor site.

Heaulme M *et al* (1986) Brain Res 384(2)
PubMedID 3022866

Sequential steps underlying neuronal plasticity induced by a transient exposure to gabazine.

Pegoraro S *et al* (2010) J Cell Physiol 222(3)
PubMedID 20027606

The kinetics of inhibition of rat recombinant heteromeric alpha1beta glycine receptors by the low-affinity antagonist SR-95531.

Beato M *et al* (2007) J Physiol 580(Pt 1)
PubMedID 17218350

The differential antagonism by bicuculline and SR95531 of pentobarbitone-induced currents in cultured hippocampal neurons.

Uchida I *et al* (1996) Eur J Pharmacol 307(1)
PubMedID 8831109

Tonically activated GABAA receptors in hippocampal neurons are high-affinity, low-conductance sensors for extracellular GABA.

Yeung et al (2003) Mol Pharmacol 36(1)
PubMedID 12488530
