


DATASHEET (S)-AMPA

Product overview

Name	(S)-AMPA
Cat No	HB0052
Biological action	Agonist
Purity	>99%
Description	AMPA receptor agonist

Images



Biological Data

Biological description

(S)-AMPA is an AMPA receptor agonist and the active enantiomer of AMPA. It is a neurotoxin in the immature rat brain.

(R,S)-AMPA is also available.

Application notes

The AMPA receptor agonist (S)-AMPA is typically used at concentrations of 1-100 μM. At 10 μM, (S)-AMPA from Hello Bio induces a large depolarising current. This depolarising current was occluded in the presence of the AMPA receptor antagonist **NBQX** (20 μM). (See Fig 1 above).

#Protocol 1: (S)-AMPA protocol

- Whole cell voltage clamp recordings of CA1 pyramidal neurons from the rat hippocampal brain slice.
- Neurons were held at -60 mV and continuously perfused with aCSF in the presence of the GABA receptor antagonist **gabazine** (20 μM).
- AMPA currents were evoked via applying (S)-AMPA directly to the recording chamber during continuous perfusion.
- To test the selectivity of (S)-AMPA to AMPA receptors, the experiment was repeated within the same neuron in the presence of the AMPA receptor antagonist **NBQX** (20 μM)
- Under these conditions (S)-AMPA failed to induce a depolarising current.

Solubility & Handling

Storage instructions

Solubility overview

Important

Room temperature

Soluble in water (100mM)

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use.

Chemical Data


Chemical name

(S)- α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid

Molecular Weight

186.17

Chemical structure



Molecular Formula

C₇H₁₀N₂O₄

CAS Number

83643-88-3

PubChem identifier

158397

SMILES

CC1=C(C(=O)NO1)C[C@@H](C(=O)O)N

Source

Synthetic

InChi

InChI=1S/C7H10N2O4/c1-3-4(6(10)9-13-3)2-5(8)7(11)12/h5H,2,8H2,1H3,(H,9,10)(H,11,12)/t5-/m0/s1

InChiKey

UUDAMDVQRQNNHZ-YFKPBYRVSA-N

MDL number

MFCD00672630

Appearance

White solid

References

Enzymic resolution and binding to rat brain membranes of the glutamic acid agonist alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid.

Hansen JJ *et al* (1983) J Med Chem 26(6)

PubMedID

[6133955](#)

Ibotenic acid analogues. Synthesis, molecular flexibility, and in vitro activity of agonists and antagonists at central glutamic acid receptors.

Lauridsen J *et al* (1985) J Med Chem 28(5)

PubMedID

[2859375](#)

The selective ionotropic-type quisqualate receptor agonist AMPA is a potent neurotoxin in immature rat brain.

McDonald JW *et al* (1990) Brain Res 526(1)

PubMedID

[1964108](#)