Hello Bio, Inc. 304 Wall St., Princeton, NJ 08540 USA

T. 609-683-7500 F. 609-228-4994

customercare-usa@m2stage.hellobio.com



DATASHEET

Streptavidin Janelia Fluor® 525

Product overview

Name Cat No Biological description Streptavidin Janelia Fluor® 525 HB15382

Streptavidin Janelia Fluor® 525 is a biotin binding protein conjugated with the fluorescent dye Janelia Fluor® 525 and can be used to detect biotin labelled molecules such as nucleic acids, antibodies, and other proteins. Biotinylated antibodies are bound with extremely high affinity by Streptavidin Janelia Fluor® 525 enabling immunofluorescent detection in IHC, ICC, flow cytometry and Western blot. Janelia Fluor® 525 and the other members of the Janelia Fluor® family are bright and highly photostable fluorophores particularly suited for super resolution imaging such as dSTORM and STED.

Key features:

- Conjugated with Janelia Fluor® 525 (Ex: 534nm, Em: 559nm)
- Supplied as a more stable lyophilate
- · Bright and photostable signal for repeated imaging
- For use in IHC(IF), ICC, Western blotting and Flow cytometry
- Suited for super resolution imaging including dSTORM and STED


Species of origin E. coli


Applications Description


fluorescence imaging, ICC, IF, IHC


Janelia Fluor® 525 conjugated streptavidin for detection and signal amplification of biotin coupled proteins and antibodies.

Images









Biological Data

Application notes

#Protocol 1: Detecting biotin-labelled antibodies in IHC

- 1. Incubate free floating rat brain sections ($40\mu m$) in sodium borohydride (NaBH₄) for 15 minutes followed by 2 hours in blocking buffer (0.05M glycine, 2% BSA and 3% donkey serum).
- 2. Incubate sections with primary antibody in blocking buffer at 4 °C overnight, as in our IHC protocol.
- 3. Wash sections three times in PBST for 5 minutes each.
- 4. Incubate sections with 2 μ g/mL goat anti-mouse biotin antibody HB11345 or goat anti-rabbit antibody HB11036 diluted in blocking buffer for 2 hours at RT.

- 5. Wash sections three times in PBST for 5 minutes each.
- 6. Incubate sections with 1 µg/mL Streptavidin Janelia Fluor® 525 in blocking buffer for 2 hours.
- 7. Wash sections three times in PBST for 5 minutes each.
- 8. Incubate sections with 10 $\mu g/mL$ DAPI for 10 minutes.
- 9. Wash sections in dH₂O, mount on glass slides with mounting media and cover with coverslip.
- 10. Image the sections on a microscope using a 532nm laser or TRITC filter set to excite Streptavidin Janelia Fluor® 525.

Solubility & Handling

Storage instructions Reconstitution advice

-20°C then use reconstitution advice