

DATASHEET

aCSF Instant Powder (Mg²⁺/Ca²⁺ free) (packets)

Product overview

Name	aCSF Instant Powder (Mg ²⁺ /Ca ²⁺ free) (packets)
Cat No	HB16218
Biological description	Artificial cerebrospinal fluid (aCSF) is a widely used buffer in electrophysiological experiments to sustain <i>ex-vivo</i> brain sections. This kit contains 20 instant powder packets. Simply dissolve the contents of each packet in dH ₂ O to a final volume of 1L, mix, add the desired concentration of Mg ²⁺ and Ca ²⁺ and bubble with carbogen to make 1L of aCSF at physiological pH. Please note: This formulation does not contain any Mg ²⁺ or Ca ²⁺ so that this can be specified by the experimenter.


Key features:

- Save time by using preformulated individual aCSF powder packets - each packet dissolves in seconds
- More reproducible with each pack's highly accurate formulation - less error for better data.
- Extensively validated in a range of patch clamp electrophysiology experiments.
- Does not contain Mg²⁺ or Ca²⁺ to allow manipulation by the experimenter. For complete aCSF containing Mg²⁺ and Ca²⁺ please see [HB9200 aCSF Instant Powder \(packets\)](#)

Contains (in mM): NaCl 124. Glucose 10, NaHCO₃ 24, KCl 3, NaH₂PO₄ 1.25

Biological action	Buffer
Description	Preformulated instant powder packets to make artificial cerebrospinal fluid (aCSF) without Mg ²⁺ or Ca ²⁺

Images



Solubility & Handling

Storage instructions RT. Dissolve each pack in dH₂O to 1L final volume.

Handling Dissolve the contents of each packet in dH₂O, add the desired quantity of Mg²⁺ and Ca²⁺ then bring the final volume to 1000ml and bubble with carbogen (10-15 minutes) to make 1L of aCSF at physiological pH. Warm to 37°C before use.

Storage instructions

RT. Dissolve each pack in dH₂O to 1L final volume.

Use immediately once opened.

Important

This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use

Chemical Data

Kit contents

Preformulated packets. Each makes 1L of aCSF.

pH after carbogenation 7.2

pH before carbogenation 7.5

References

Reduced expression of the psychiatric risk gene DLG2 (PSD93) impairs hippocampal synaptic integration and plasticity.

Griesius S et al (2022) Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 47

PubMedID [35115661](#)

The development of synaptic plasticity induction rules and the requirement for postsynaptic spikes in rat hippocampal CA1 pyramidal neurones.

Buchanan KA et al (2007) The Journal of physiology 585

PubMedID [17932146](#)
